

IMO Shortlist 2014

www.artofproblemsolving.com/community/c107000

by Sayan, Amir Hossein, hajimbrak, v_Enhance, IMO2018, ipaper, codyj

– Algebra

- A1** Let $a_0 < a_1 < a_2 < \dots$ be an infinite sequence of positive integers. Prove that there exists a unique integer $n \geq 1$ such that

$$a_n < \frac{a_0 + a_1 + a_2 + \dots + a_n}{n} \leq a_{n+1}.$$

Proposed by Gerhard Wöginger, Austria.

- A2** Define the function $f : (0, 1) \rightarrow (0, 1)$ by

$$f(x) = \begin{cases} x + \frac{1}{2} & \text{if } x < \frac{1}{2} \\ x^2 & \text{if } x \geq \frac{1}{2} \end{cases}$$

Let a and b be two real numbers such that $0 < a < b < 1$. We define the sequences a_n and b_n by $a_0 = a, b_0 = b$, and $a_n = f(a_{n-1}), b_n = f(b_{n-1})$ for $n > 0$. Show that there exists a positive integer n such that

$$(a_n - a_{n-1})(b_n - b_{n-1}) < 0.$$

Proposed by Denmark

- A3** For a sequence x_1, x_2, \dots, x_n of real numbers, we define its *price* as

$$\max_{1 \leq i \leq n} |x_1 + \dots + x_i|.$$

Given n real numbers, Dave and George want to arrange them into a sequence with a low price. Diligent Dave checks all possible ways and finds the minimum possible price D . Greedy George, on the other hand, chooses x_1 such that $|x_1|$ is as small as possible; among the remaining numbers, he chooses x_2 such that $|x_1 + x_2|$ is as small as possible, and so on. Thus, in the i -th step he chooses x_i among the remaining numbers so as to minimise the value of $|x_1 + x_2 + \dots + x_i|$. In each step, if several numbers provide the same value, George chooses one at random. Finally he gets a sequence with price G .

Find the least possible constant c such that for every positive integer n , for every collection of n real numbers, and for every possible sequence that George might obtain, the resulting values satisfy the inequality $G \leq cD$.

Proposed by Georgia

- A4** Determine all functions $f : \mathbb{Z} \rightarrow \mathbb{Z}$ satisfying

$$f(f(m) + n) + f(m) = f(n) + f(3m) + 2014$$

for all integers m and n .

Proposed by Netherlands

- A5** Consider all polynomials $P(x)$ with real coefficients that have the following property: for any two real numbers x and y one has

$$|y^2 - P(x)| \leq 2|x| \quad \text{if and only if} \quad |x^2 - P(y)| \leq 2|y|.$$

Determine all possible values of $P(0)$.

Proposed by Belgium

- A6** Find all functions $f : \mathbb{Z} \rightarrow \mathbb{Z}$ such that

$$n^2 + 4f(n) = f(f(n))^2$$

for all $n \in \mathbb{Z}$.

Proposed by Sahl Khan, UK

– Combinatorics

- C1** Let n points be given inside a rectangle R such that no two of them lie on a line parallel to one of the sides of R . The rectangle R is to be dissected into smaller rectangles with sides parallel to the sides of R in such a way that none of these rectangles contains any of the given points in its interior. Prove that we have to dissect R into at least $n + 1$ smaller rectangles.

Proposed by Serbia

- C2** We have 2^m sheets of paper, with the number 1 written on each of them. We perform the following operation. In every step we choose two distinct sheets; if the numbers on the two sheets are a and b , then we erase these numbers and write the number $a + b$ on both sheets. Prove that after $m2^{m-1}$ steps, the sum of the numbers on all the sheets is at least 4^m .

Proposed by Abbas Mehrabian, Iran

- C3** Let $n \geq 2$ be an integer. Consider an $n \times n$ chessboard consisting of n^2 unit squares. A configuration of n rooks on this board is *peaceful* if every row and every column contains exactly one rook. Find the greatest positive integer k such that, for each peaceful configuration of n rooks, there is a $k \times k$ square which does not contain a rook on any of its k^2 unit squares.
-

- C4** Construct a tetromino by attaching two 2×1 dominoes along their longer sides such that the midpoint of the longer side of one domino is a corner of the other domino. This construction yields two kinds of tetrominoes with opposite orientations. Let us call them S - and Z -tetrominoes, respectively.

Assume that a lattice polygon P can be tiled with S -tetrominoes. Prove that no matter how we tile P using only S - and Z -tetrominoes, we always use an even number of Z -tetrominoes.

Proposed by Tamas Fleiner and Peter Pal Pach, Hungary

- C5** A set of lines in the plane is in *general position* if no two are parallel and no three pass through the same point. A set of lines in general position cuts the plane into regions, some of which have finite area; we call these its *finite regions*. Prove that for all sufficiently large n , in any set of n lines in general position it is possible to colour at least \sqrt{n} lines blue in such a way that none of its finite regions has a completely blue boundary.

Note: Results with \sqrt{n} replaced by $c\sqrt{n}$ will be awarded points depending on the value of the constant c .

- C6** We are given an infinite deck of cards, each with a real number on it. For every real number x , there is exactly one card in the deck that has x written on it. Now two players draw disjoint sets A and B of 100 cards each from this deck. We would like to define a rule that declares one of them a winner. This rule should satisfy the following conditions:

1. The winner only depends on the relative order of the 200 cards: if the cards are laid down in increasing order face down and we are told which card belongs to which player, but not what numbers are written on them, we can still decide the winner.

2. If we write the elements of both sets in increasing order as $A = \{a_1, a_2, \dots, a_{100}\}$ and $B = \{b_1, b_2, \dots, b_{100}\}$, and $a_i > b_i$ for all i , then A beats B .

3. If three players draw three disjoint sets A, B, C from the deck, A beats B and B beats C then A also beats C .

How many ways are there to define such a rule? Here, we consider two rules as different if there exist two sets A and B such that A beats B according to one rule, but B beats A according to the other.

Proposed by Ilya Bogdanov, Russia

- C7** Let M be a set of $n \geq 4$ points in the plane, no three of which are collinear. Initially these points are connected with n segments so that each point in M is the endpoint of exactly two segments. Then, at each step, one may choose two segments AB and CD sharing a common interior point and replace them by the segments AC and BD if none of them is present at this moment. Prove that it is impossible to perform $n^3/4$ or more such moves.

Proposed by Vladislav Volkov, Russia

- C8** A card deck consists of 1024 cards. On each card, a set of distinct decimal digits is written in such a way that no two of these sets coincide (thus, one of the cards is empty). Two players

alternately take cards from the deck, one card per turn. After the deck is empty, each player checks if he can throw out one of his cards so that each of the ten digits occurs on an even number of his remaining cards. If one player can do this but the other one cannot, the one who can is the winner; otherwise a draw is declared.

Determine all possible first moves of the first player after which he has a winning strategy.

Proposed by Ilya Bogdanov & Vladimir Bragin, Russia

- C9** There are n circles drawn on a piece of paper in such a way that any two circles intersect in two points, and no three circles pass through the same point. Turbo the snail slides along the circles in the following fashion. Initially he moves on one of the circles in clockwise direction. Turbo always keeps sliding along the current circle until he reaches an intersection with another circle. Then he continues his journey on this new circle and also changes the direction of moving, i.e. from clockwise to anticlockwise or *vice versa*. Suppose that Turbo's path entirely covers all circles. Prove that n must be odd.

Proposed by Tejaswi Navilarekallu, India

– Geometry

- G1** Let P and Q be on segment BC of an acute triangle ABC such that $\angle PAB = \angle BCA$ and $\angle CAQ = \angle ABC$. Let M and N be the points on AP and AQ , respectively, such that P is the midpoint of AM and Q is the midpoint of AN . Prove that the intersection of BM and CN is on the circumference of triangle ABC .

Proposed by Giorgi Arabidze, Georgia.

- G2** Let ABC be a triangle. The points K, L , and M lie on the segments BC, CA , and AB , respectively, such that the lines AK, BL , and CM intersect in a common point. Prove that it is possible to choose two of the triangles ALM, BMK , and CKL whose inradii sum up to at least the inradius of the triangle ABC .

Proposed by Estonia

- G3** Let Ω and O be the circumcircle and the circumcentre of an acute-angled triangle ABC with $AB > BC$. The angle bisector of $\angle ABC$ intersects Ω at $M \neq B$. Let Γ be the circle with diameter BM . The angle bisectors of $\angle AOB$ and $\angle BOC$ intersect Γ at points P and Q , respectively. The point R is chosen on the line PQ so that $BR = MR$. Prove that $BR \parallel AC$. (Here we always assume that an angle bisector is a ray.)

Proposed by Sergey Berlov, Russia

- G4** Consider a fixed circle Γ with three fixed points A, B , and C on it. Also, let us fix a real number $\lambda \in (0, 1)$. For a variable point $P \notin \{A, B, C\}$ on Γ , let M be the point on the segment CP such

that $CM = \lambda \cdot CP$. Let Q be the second point of intersection of the circumcircles of the triangles AMP and BMC . Prove that as P varies, the point Q lies on a fixed circle.

Proposed by Jack Edward Smith, UK

- G5** Convex quadrilateral $ABCD$ has $\angle ABC = \angle CDA = 90^\circ$. Point H is the foot of the perpendicular from A to BD . Points S and T lie on sides AB and AD , respectively, such that H lies inside triangle SCT and

$$\angle CHS - \angle CSB = 90^\circ, \quad \angle THC - \angle DTC = 90^\circ.$$

Prove that line BD is tangent to the circumcircle of triangle TSH .

- G6** Let ABC be a fixed acute-angled triangle. Consider some points E and F lying on the sides AC and AB , respectively, and let M be the midpoint of EF . Let the perpendicular bisector of EF intersect the line BC at K , and let the perpendicular bisector of MK intersect the lines AC and AB at S and T , respectively. We call the pair (E, F) *interesting*, if the quadrilateral $KSAT$ is cyclic.

Suppose that the pairs (E_1, F_1) and (E_2, F_2) are interesting. Prove that $\frac{E_1E_2}{AB} = \frac{F_1F_2}{AC}$

Proposed by Ali Zamani, Iran

- G7** Let ABC be a triangle with circumcircle Ω and incentre I . Let the line passing through I and perpendicular to CI intersect the segment BC and the arc BC (not containing A) of Ω at points U and V , respectively. Let the line passing through U and parallel to AI intersect AV at X , and let the line passing through V and parallel to AI intersect AB at Y . Let W and Z be the midpoints of AX and BC , respectively. Prove that if the points I, X , and Y are collinear, then the points I, W , and Z are also collinear.

Proposed by David B. Rush, USA

– Number Theory

- N1** Let $n \geq 2$ be an integer, and let A_n be the set

$$A_n = \{2^n - 2^k \mid k \in \mathbb{Z}, 0 \leq k < n\}.$$

Determine the largest positive integer that cannot be written as the sum of one or more (not necessarily distinct) elements of A_n .

Proposed by Serbia

- N2** Determine all pairs (x, y) of positive integers such that

$$\sqrt[3]{7x^2 - 13xy + 7y^2} = |x - y| + 1.$$

Proposed by Titu Andreescu, USA

- N3** For each positive integer n , the Bank of Cape Town issues coins of denomination $\frac{1}{n}$. Given a finite collection of such coins (of not necessarily different denominations) with total value at most $99 + \frac{1}{2}$, prove that it is possible to split this collection into 100 or fewer groups, such that each group has total value at most 1.

- N4** Let $n > 1$ be a given integer. Prove that infinitely many terms of the sequence $(a_k)_{k \geq 1}$, defined by

$$a_k = \left\lfloor \frac{n^k}{k} \right\rfloor,$$

are odd. (For a real number x , $\lfloor x \rfloor$ denotes the largest integer not exceeding x .)

Proposed by Hong Kong

- N5** Find all triples (p, x, y) consisting of a prime number p and two positive integers x and y such that $x^{p-1} + y$ and $x + y^{p-1}$ are both powers of p .

Proposed by Belgium

- N6** Let $a_1 < a_2 < \dots < a_n$ be pairwise coprime positive integers with a_1 being prime and $a_1 \geq n + 2$. On the segment $I = [0, a_1 a_2 \dots a_n]$ of the real line, mark all integers that are divisible by at least one of the numbers a_1, \dots, a_n . These points split I into a number of smaller segments. Prove that the sum of the squares of the lengths of these segments is divisible by a_1 .

Proposed by Serbia

- N7** Let $c \geq 1$ be an integer. Define a sequence of positive integers by $a_1 = c$ and

$$a_{n+1} = a_n^3 - 4c \cdot a_n^2 + 5c^2 \cdot a_n + c$$

for all $n \geq 1$. Prove that for each integer $n \geq 2$ there exists a prime number p dividing a_n but none of the numbers a_1, \dots, a_{n-1} .

Proposed by Austria

- N8** For every real number x , let $\|x\|$ denote the distance between x and the nearest integer. Prove that for every pair (a, b) of positive integers there exist an odd prime p and a positive integer k satisfying

$$\left\| \frac{a}{p^k} \right\| + \left\| \frac{b}{p^k} \right\| + \left\| \frac{a+b}{p^k} \right\| = 1.$$

Proposed by Geza Kos, Hungary