

IMO Shortlist 2015

www.artofproblemsolving.com/community/c111148

by gavrilos, ABCDE, va2010, codyj, CantonMathGuy, randomusername, Problem_Penetrator, samithay-ohan, termas

– Algebra

A1 Suppose that a sequence a_1, a_2, \dots of positive real numbers satisfies

$$a_{k+1} \geq \frac{ka_k}{a_k^2 + (k-1)}$$

for every positive integer k . Prove that $a_1 + a_2 + \dots + a_n \geq n$ for every $n \geq 2$.

A2 Determine all functions $f : \mathbb{Z} \rightarrow \mathbb{Z}$ with the property that

$$f(x - f(y)) = f(f(x)) - f(y) - 1$$

holds for all $x, y \in \mathbb{Z}$.

A3 Let n be a fixed positive integer. Find the maximum possible value of

$$\sum_{1 \leq r < s \leq 2n} (s - r - n)x_r x_s,$$

where $-1 \leq x_i \leq 1$ for all $i = 1, \dots, 2n$.

A4 Let \mathbb{R} be the set of real numbers. Determine all functions $f : \mathbb{R} \rightarrow \mathbb{R}$ that satisfy the equation

$$f(x + f(x + y)) + f(xy) = x + f(x + y) + yf(x)$$

for all real numbers x and y .

Proposed by Dorlir Ahmeti, Albania

A5 Let $2\mathbb{Z} + 1$ denote the set of odd integers. Find all functions $f : \mathbb{Z} \mapsto 2\mathbb{Z} + 1$ satisfying

$$f(x + f(x) + y) + f(x - f(x) - y) = f(x + y) + f(x - y)$$

for every $x, y \in \mathbb{Z}$.

A6 Let n be a fixed integer with $n \geq 2$. We say that two polynomials P and Q with real coefficients are *block-similar* if for each $i \in \{1, 2, \dots, n\}$ the sequences

$$P(2015i), P(2015i - 1), \dots, P(2015i - 2014) \quad \text{and} \\ Q(2015i), Q(2015i - 1), \dots, Q(2015i - 2014)$$

are permutations of each other.

- (a) Prove that there exist distinct block-similar polynomials of degree $n + 1$.
 (b) Prove that there do not exist distinct block-similar polynomials of degree n .

Proposed by David Arthur, Canada

– Combinatorics

- C1** In Lineland there are $n \geq 1$ towns, arranged along a road running from left to right. Each town has a *left bulldozer* (put to the left of the town and facing left) and a *right bulldozer* (put to the right of the town and facing right). The sizes of the $2n$ bulldozers are distinct. Every time when a left and right bulldozer confront each other, the larger bulldozer pushes the smaller one off the road. On the other hand, bulldozers are quite unprotected at their rears; so, if a bulldozer reaches the rear-end of another one, the first one pushes the second one off the road, regardless of their sizes.

Let A and B be two towns, with B to the right of A . We say that town A can *sweep* town B away if the right bulldozer of A can move over to B pushing off all bulldozers it meets. Similarly town B can sweep town A away if the left bulldozer of B can move over to A pushing off all bulldozers of all towns on its way.

Prove that there is exactly one town that cannot be swept away by any other one.

- C2** We say that a finite set S of points in the plane is *balanced* if, for any two different points A and B in S , there is a point C in S such that $AC = BC$. We say that S is *centre-free* if for any three different points A, B and C in S , there is no points P in S such that $PA = PB = PC$.

- (a) Show that for all integers $n \geq 3$, there exists a balanced set consisting of n points.
 (b) Determine all integers $n \geq 3$ for which there exists a balanced centre-free set consisting of n points.

Proposed by Netherlands

- C3** For a finite set A of positive integers, a partition of A into two disjoint nonempty subsets A_1 and A_2 is *good* if the least common multiple of the elements in A_1 is equal to the greatest common divisor of the elements in A_2 . Determine the minimum value of n such that there exists a set of n positive integers with exactly 2015 good partitions.

- C4** Let n be a positive integer. Two players A and B play a game in which they take turns choosing positive integers $k \leq n$. The rules of the game are:
- (i) A player cannot choose a number that has been chosen by either player on any previous turn.
 - (ii) A player cannot choose a number consecutive to any of those the player has already chosen on any previous turn.
 - (iii) The game is a draw if all numbers have been chosen; otherwise the player who cannot choose a number anymore loses the game.

The player A takes the first turn. Determine the outcome of the game, assuming that both players use optimal strategies.

Proposed by Finland

- C5** The sequence a_1, a_2, \dots of integers satisfies the conditions:

- (i) $1 \leq a_j \leq 2015$ for all $j \geq 1$,
- (ii) $k + a_k \neq \ell + a_\ell$ for all $1 \leq k < \ell$.

Prove that there exist two positive integers b and N for which

$$\left| \sum_{j=m+1}^n (a_j - b) \right| \leq 1007^2$$

for all integers m and n such that $n > m \geq N$.

Proposed by Ivan Guo and Ross Atkins, Australia

- C6** Let S be a nonempty set of positive integers. We say that a positive integer n is *clean* if it has a unique representation as a sum of an odd number of distinct elements from S . Prove that there exist infinitely many positive integers that are not clean.

- C7** In a company of people some pairs are enemies. A group of people is called *unsociable* if the number of members in the group is odd and at least 3, and it is possible to arrange all its members around a round table so that every two neighbors are enemies. Given that there are at most 2015 unsociable groups, prove that it is possible to partition the company into 11 parts so that no two enemies are in the same part.

Proposed by Russia

– Geometry

- G1** Let ABC be an acute triangle with orthocenter H . Let G be the point such that the quadrilateral $ABGH$ is a parallelogram. Let I be the point on the line GH such that AC bisects HI . Suppose

that the line AC intersects the circumcircle of the triangle GCI at C and J . Prove that $IJ = AH$.

- G2** Triangle ABC has circumcircle Ω and circumcenter O . A circle Γ with center A intersects the segment BC at points D and E , such that B, D, E , and C are all different and lie on line BC in this order. Let F and G be the points of intersection of Γ and Ω , such that A, F, B, C , and G lie on Ω in this order. Let K be the second point of intersection of the circumcircle of triangle BDF and the segment AB . Let L be the second point of intersection of the circumcircle of triangle CGE and the segment CA .

Suppose that the lines FK and GL are different and intersect at the point X . Prove that X lies on the line AO .

Proposed by Greece

- G3** Let ABC be a triangle with $\angle C = 90^\circ$, and let H be the foot of the altitude from C . A point D is chosen inside the triangle CBH so that CH bisects AD . Let P be the intersection point of the lines BD and CH . Let ω be the semicircle with diameter BD that meets the segment CB at an interior point. A line through P is tangent to ω at Q . Prove that the lines CQ and AD meet on ω .

- G4** Let ABC be an acute triangle and let M be the midpoint of AC . A circle ω passing through B and M meets the sides AB and BC at points P and Q respectively. Let T be the point such that $BPTQ$ is a parallelogram. Suppose that T lies on the circumcircle of ABC . Determine all possible values of $\frac{BT}{BM}$.

- G5** Let ABC be a triangle with $CA \neq CB$. Let D, F , and G be the midpoints of the sides AB, AC , and BC respectively. A circle Γ passing through C and tangent to AB at D meets the segments AF and BG at H and I , respectively. The points H' and I' are symmetric to H and I about F and G , respectively. The line $H'I'$ meets CD and FG at Q and M , respectively. The line CM meets Γ again at P . Prove that $CQ = QP$.

Proposed by El Salvador

- G6** Let ABC be an acute triangle with $AB > AC$. Let Γ be its circumcircle, H its orthocenter, and F the foot of the altitude from A . Let M be the midpoint of BC . Let Q be the point on Γ such that $\angle HQA = 90^\circ$ and let K be the point on Γ such that $\angle HKQ = 90^\circ$. Assume that the points A, B, C, K and Q are all different and lie on Γ in this order.

Prove that the circumcircles of triangles KQH and FKM are tangent to each other.

Proposed by Ukraine

- G7** Let $ABCD$ be a convex quadrilateral, and let P, Q, R , and S be points on the sides AB, BC, CD , and DA , respectively. Let the line segment PR and QS meet at O . Suppose that each of

the quadrilaterals $APOS$, $BQOP$, $CROQ$, and $DSOR$ has an incircle. Prove that the lines AC , PQ , and RS are either concurrent or parallel to each other.

- G8** A *triangulation* of a convex polygon Π is a partitioning of Π into triangles by diagonals having no common points other than the vertices of the polygon. We say that a triangulation is a *Thaiangulation* if all triangles in it have the same area.

Prove that any two different Thaiangulations of a convex polygon Π differ by exactly two triangles. (In other words, prove that it is possible to replace one pair of triangles in the first Thaiangulation with a different pair of triangles so as to obtain the second Thaiangulation.)

Proposed by Bulgaria

- Number Theory

- N1** Determine all positive integers M such that the sequence a_0, a_1, a_2, \dots defined by

$$a_0 = M + \frac{1}{2} \quad \text{and} \quad a_{k+1} = a_k \lfloor a_k \rfloor \quad \text{for } k = 0, 1, 2, \dots$$

contains at least one integer term.

- N2** Let a and b be positive integers such that $a! + b!$ divides $a!b!$. Prove that $3a \geq 2b + 2$.

- N3** Let m and n be positive integers such that $m > n$. Define $x_k = \frac{m+k}{n+k}$ for $k = 1, 2, \dots, n+1$. Prove that if all the numbers x_1, x_2, \dots, x_{n+1} are integers, then $x_1 x_2 \dots x_{n+1} - 1$ is divisible by an odd prime.

- N4** Suppose that a_0, a_1, \dots and b_0, b_1, \dots are two sequences of positive integers such that $a_0, b_0 \geq 2$ and

$$a_{n+1} = \gcd(a_n, b_n) + 1, \quad b_{n+1} = \text{lcm}(a_n, b_n) - 1.$$

Show that the sequence a_n is eventually periodic; in other words, there exist integers $N \geq 0$ and $t > 0$ such that $a_{n+t} = a_n$ for all $n \geq N$.

- N5** Find all positive integers (a, b, c) such that

$$ab - c, \quad bc - a, \quad ca - b$$

are all powers of 2.

Proposed by Serbia

- N6** Let $\mathbb{Z}_{>0}$ denote the set of positive integers. Consider a function $f : \mathbb{Z}_{>0} \rightarrow \mathbb{Z}_{>0}$. For any $m, n \in \mathbb{Z}_{>0}$ we write $f^n(m) = \underbrace{f(f(\dots f(m)\dots))}_n$. Suppose that f has the following two properties:

- (i) if $m, n \in \mathbb{Z}_{>0}$, then $\frac{f^n(m)-m}{n} \in \mathbb{Z}_{>0}$;
 (ii) The set $\mathbb{Z}_{>0} \setminus \{f(n) \mid n \in \mathbb{Z}_{>0}\}$ is finite.

Prove that the sequence $f(1) - 1, f(2) - 2, f(3) - 3, \dots$ is periodic.

Proposed by Ang Jie Jun, Singapore

- N7** Let $\mathbb{Z}_{>0}$ denote the set of positive integers. For any positive integer k , a function $f : \mathbb{Z}_{>0} \rightarrow \mathbb{Z}_{>0}$ is called **[i] k -good** if $\gcd(f(m) + n, f(n) + m) \leq k$ for all $m \neq n$. Find all k such that there exists a k -good function.

Proposed by James Rickards, Canada

- N8** For every positive integer n with prime factorization $n = \prod_{i=1}^k p_i^{\alpha_i}$, define

$$\mathcal{U}(n) = \sum_{i: p_i > 10^{100}} \alpha_i.$$

That is, $\mathcal{U}(n)$ is the number of prime factors of n greater than 10^{100} , counted with multiplicity.

Find all strictly increasing functions $f : \mathbb{Z} \rightarrow \mathbb{Z}$ such that

$$\mathcal{U}(f(a) - f(b)) \leq \mathcal{U}(a - b) \quad \text{for all integers } a \text{ and } b \text{ with } a > b.$$

Proposed by Rodrigo Sanches Angelo, Brazil