

Rice Math Tournament 2006
www.artofproblemsolving.com/community/c2976668

by parmenides51, phiReKaLk6781, bluecarneal

– Team Round

-
- 1** Given $\triangle ABC$, where A is at $(0, 0)$, B is at $(20, 0)$, and C is on the positive y -axis. Cone M is formed when $\triangle ABC$ is rotated about the x -axis, and cone N is formed when $\triangle ABC$ is rotated about the y -axis. If the volume of cone M minus the volume of cone N is 140π , find the length of \overline{BC} .
-
- 2** In a given sequence $\{S_1, S_2, \dots, S_k\}$, for terms $n \geq 3$, $S_n = \sum_{i=1}^{n-1} i \cdot S_{n-i}$. For example, if the first two elements are 2 and 3, respectively, the third entry would be $1 \cdot 3 + 2 \cdot 2 = 7$, and the fourth would be $1 \cdot 7 + 2 \cdot 3 + 3 \cdot 2 = 19$, and so on. Given that a sequence of integers having this form starts with 2, and the 7th element is 68, what is the second element?
-
- 3** A triangle has altitudes of length 5 and 7. What is the maximum length of the third altitude?
-
- 4** Let $x + y = a$ and $xy = b$. The expression $x^6 + y^6$ can be written as a polynomial in terms of a and b . What is this polynomial?
-
- 5** There exist two positive numbers x such that $\sin(\arccos(\tan(\arcsin x))) = x$. Find the product of the two possible x .
-
- 6** The expression $16^n + 4^n + 1$ is equivalent to the expression $(2^{p(n)} - 1)/(2^{q(n)} - 1)$ for all positive integers $n > 1$ where $p(n)$ and $q(n)$ are functions and $\frac{p(n)}{q(n)}$ is constant. Find $p(2006) - q(2006)$.
-
- 7** Let S be the set of all 3-tuples (a, b, c) that satisfy $a + b + c = 3000$ and $a, b, c > 0$. If one of these 3-tuples is chosen at random, what's the probability that a, b or c is greater than or equal to 2,500?
-
- 8** Evaluate: $\lim_{n \rightarrow \infty} \sum_{k=n^2}^{(n+1)^2} \frac{1}{\sqrt{k}}$
-
- 9** $\triangle ABC$ has $AB = AC$. Points M and N are midpoints of \overline{AB} and \overline{AC} , respectively. The medians \overline{MC} and \overline{NB} intersect at a right angle. Find $(\frac{AB}{BC})^2$.
-
- 10** Find the smallest positive m for which there are at least 11 even and 11 odd positive integers n so that $\frac{n^3+m}{n+2}$ is an integer.
-

11 Polynomial $P(x) = c_{2006}x^{2006} + c_{2005}x^{2005} + \dots + c_1x + c_0$ has roots $r_1, r_2, \dots, r_{2006}$. The coefficients satisfy $2i \frac{c_i}{c_{2006-i}} = 2j \frac{c_j}{c_{2006-j}}$ for all pairs of integers $0 \leq i, j \leq 2006$. Given that $\sum_{i \neq j, i=1, j=1}^{2006} \frac{r_i}{r_j} = 42$, determine $\sum_{i=1}^{2006} (r_1 + r_2 + \dots + r_{2006})$.

12 Find the total number of k -tuples (n_1, n_2, \dots, n_k) of positive integers so that $n_{i+1} \geq n_i$ for each i , and k regular polygons with numbers of sides n_1, n_2, \dots, n_k respectively will fit into a tessellation at a point. That is, the sum of one interior angle from each of the polygons is 360° .

13 A ray is drawn from the origin tangent to the graph of the upper part of the hyperbola $y^2 = x^2 - x + 1$ in the first quadrant. This ray makes an angle of θ with the positive x -axis. Compute $\cos \theta$.

14 Find the smallest nonnegative integer n for which $\binom{2006}{n}$ is divisible by 7^3 .

15 Let c_i denote the i th composite integer so that $\{c_i\} = 4, 6, 8, 9, \dots$. Compute

$$\prod_{i=1}^{\infty} \frac{c_i^2}{c_i^2 - 1}$$

(Hint: $\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$)