

Romanian Masters In Mathematics 2010

www.artofproblemsolving.com/community/c3609

by Goutham

- 1 For a finite non empty set of primes P , let $m(P)$ denote the largest possible number of consecutive positive integers, each of which is divisible by at least one member of P .

(i) Show that $|P| \leq m(P)$, with equality if and only if $\min(P) > |P|$.

(ii) Show that $m(P) < (|P| + 1)(2^{|P|} - 1)$.

(The number $|P|$ is the size of set P)

Dan Schwarz, Romania

- 2 For each positive integer n , find the largest real number C_n with the following property. Given any n real-valued functions $f_1(x), f_2(x), \dots, f_n(x)$ defined on the closed interval $0 \leq x \leq 1$, one can find numbers x_1, x_2, \dots, x_n , such that $0 \leq x_i \leq 1$ satisfying

$$|f_1(x_1) + f_2(x_2) + \dots + f_n(x_n) - x_1 x_2 \dots x_n| \geq C_n$$

Marko Radovanovi, Serbia

- 3 Let $A_1 A_2 A_3 A_4$ be a quadrilateral with no pair of parallel sides. For each $i = 1, 2, 3, 4$, define ω_i to be the circle touching the quadrilateral externally, and which is tangent to the lines $A_{i-1} A_i, A_i A_{i+1}$ and $A_{i+1} A_{i+2}$ (indices are considered modulo 4 so $A_0 = A_4, A_5 = A_1$ and $A_6 = A_2$). Let T_i be the point of tangency of ω_i with the side $A_i A_{i+1}$. Prove that the lines $A_1 A_2, A_3 A_4$ and $T_2 T_4$ are concurrent if and only if the lines $A_2 A_3, A_4 A_1$ and $T_1 T_3$ are concurrent.

Pavel Kozhevnikov, Russia

- 4 Determine whether there exists a polynomial $f(x_1, x_2)$ with two variables, with integer coefficients, and two points $A = (a_1, a_2)$ and $B = (b_1, b_2)$ in the plane, satisfying the following conditions:

(i) A is an integer point (i.e a_1 and a_2 are integers);

(ii) $|a_1 - b_1| + |a_2 - b_2| = 2010$;

(iii) $f(n_1, n_2) > f(a_1, a_2)$ for all integer points (n_1, n_2) in the plane other than A ;

(iv) $f(x_1, x_2) > f(b_1, b_2)$ for all integer points (x_1, x_2) in the plane other than B .

Massimo Gobbino, Italy

- 5 Let n be a given positive integer. Say that a set K of points with integer coordinates in the plane is connected if for every pair of points $R, S \in K$, there exists a positive integer ℓ and a sequence $R = T_0, T_1, T_2, \dots, T_\ell = S$ of points in K , where each T_i is distance 1 away from T_{i+1} . For such a set K , we define the set of vectors

$$\Delta(K) = \{\overrightarrow{RS} \mid R, S \in K\}$$

What is the maximum value of $|\Delta(K)|$ over all connected sets K of $2n + 1$ points with integer coordinates in the plane?

Grigory Chelnokov, Russia

-
- 6 Given a polynomial $f(x)$ with rational coefficients, of degree $d \geq 2$, we define the sequence of sets $f^0(\mathbb{Q}), f^1(\mathbb{Q}), \dots$ as $f^0(\mathbb{Q}) = \mathbb{Q}$, $f^{n+1}(\mathbb{Q}) = f(f^n(\mathbb{Q}))$ for $n \geq 0$. (Given a set S , we write $f(S)$ for the set $\{f(x) \mid x \in S\}$). Let $f^\omega(\mathbb{Q}) = \bigcap_{n=0}^{\infty} f^n(\mathbb{Q})$ be the set of numbers that are in all of the sets $f^n(\mathbb{Q})$, $n \geq 0$. Prove that $f^\omega(\mathbb{Q})$ is a finite set.

Dan Schwarz, Romania
