2012 Balkan MO



AoPS Community

Balkan MO 2012

www.artofproblemsolving.com/community/c4084 by cefer

-	April 28th
-	April 28th

- 1 Let *A*, *B* and *C* be points lying on a circle Γ with centre *O*. Assume that $\angle ABC > 90$. Let *D* be the point of intersection of the line *AB* with the line perpendicular to *AC* at *C*. Let *l* be the line through *D* which is perpendicular to *AO*. Let *E* be the point of intersection of *l* with the line *AC*, and let *F* be the point of intersection of Γ with *l* that lies between *D* and *E*. Prove that the circumcircles of triangles *BFE* and *CFD* are tangent at *F*.
- **2** Prove that

$$\sum_{cyc} (x+y)\sqrt{(z+x)(z+y)} \ge 4(xy+yz+zx),$$

for all positive real numbers x, y and z.

- **3** Let *n* be a positive integer. Let $P_n = \{2^n, 2^{n-1} \cdot 3, 2^{n-2} \cdot 3^2, \dots, 3^n\}$. For each subset *X* of P_n , we write S_X for the sum of all elements of *X*, with the convention that $S_{\emptyset} = 0$ where \emptyset is the empty set. Suppose that *y* is a real number with $0 \le y \le 3^{n+1} 2^{n+1}$. Prove that there is a subset *Y* of P_n such that $0 \le y S_Y < 2^n$
- **4** Let \mathbb{Z}^+ be the set of positive integers. Find all functions $f : \mathbb{Z}^+ \to \mathbb{Z}^+$ such that the following conditions both hold:

(i) f(n!) = f(n)! for every positive integer n,

(ii) m - n divides f(m) - f(n) whenever m and n are different positive integers.

Art of Problem Solving is an ACS WASC Accredited School.