

ELMO Shortlist 2011

www.artofproblemsolving.com/community/c4349

by math154

– Algebra

- 1** Let n be a positive integer. There are n soldiers stationed on the n th root of unity in the complex plane. Each round, you pick a point, and all the soldiers shoot in a straight line towards that point; if their shot hits another soldier, the hit soldier dies and no longer shoots during the next round. What is the minimum number of rounds, in terms of n , required to eliminate all the soldiers?

David Yang.

- 2** Find all functions $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that whenever $a > b > c > d > 0$ and $ad = bc$,

$$f(a + d) + f(b - c) = f(a - d) + f(b + c).$$

Calvin Deng.

- 3** Let N be a positive integer. Define a sequence a_0, a_1, \dots by $a_0 = 0$, $a_1 = 1$, and $a_{n+1} + a_{n-1} = a_n(2 - 1/N)$ for $n \geq 1$. Prove that $a_n < \sqrt{N + 1}$ for all n .

Evan O'Dorney.

- 4** In terms of $n \geq 2$, find the largest constant c such that for all nonnegative a_1, a_2, \dots, a_n satisfying $a_1 + a_2 + \dots + a_n = n$, the following inequality holds:

$$\frac{1}{n + ca_1^2} + \frac{1}{n + ca_2^2} + \dots + \frac{1}{n + ca_n^2} \leq \frac{n}{n + c}.$$

Calvin Deng.

- 5** Given positive reals x, y, z such that $xy + yz + zx = 1$, show that

$$\sum_{\text{cyc}} \sqrt{(xy + kx + ky)(xz + kx + kz)} \geq k^2,$$

where $k = 2 + \sqrt{3}$.

Victor Wang.

- 6 Let $Q(x)$ be a polynomial with integer coefficients. Prove that there exists a polynomial $P(x)$ with integer coefficients such that for every integer $n \geq \deg Q$,

$$\sum_{i=0}^n \frac{!i P(i)}{i!(n-i)!} = Q(n),$$

where $!i$ denotes the number of derangements (permutations with no fixed points) of $1, 2, \dots, i$.

Calvin Deng.

- 7 Determine whether there exist two reals x, y and a sequence $\{a_n\}_{n=0}^{\infty}$ of nonzero reals such that $a_{n+2} = xa_{n+1} + ya_n$ for all $n \geq 0$ and for every positive real number r , there exist positive integers i, j such that $|a_i| < r < |a_j|$.

Alex Zhu.

- 8 Let $n > 1$ be an integer and a, b, c be three complex numbers such that $a + b + c = 0$ and $a^n + b^n + c^n = 0$. Prove that two of a, b, c have the same magnitude.

Evan O'Dorney.

– Combinatorics

- 1 Let S be a finite set, and let F be a family of subsets of S such that

a) If $A \subseteq S$, then $A \in F$ if and only if $S \setminus A \notin F$;

b) If $A \subseteq B \subseteq S$ and $B \in F$, then $A \in F$.

Determine if there must exist a function $f : S \rightarrow \mathbb{R}$ such that for every $A \subseteq S$, $A \in F$ if and only if

$$\sum_{s \in A} f(s) < \sum_{s \in S \setminus A} f(s).$$

Evan O'Dorney.

- 2 A directed graph has each vertex with outdegree 2. Prove that it is possible to split the vertices into 3 sets so that for each vertex v , v is not simultaneously in the same set with both of the vertices that it points to.

David Yang.

See here (<http://www.artofproblemsolving.com/Forum/viewtopic.php?f=42&t=492100>).

- 3 Wanda the Worm likes to eat Pascal's triangle. One day, she starts at the top of the triangle and eats $\binom{0}{0} = 1$. Each move, she travels to an adjacent positive integer and eats it, but she can never return to a spot that she has previously eaten. If Wanda can never eat numbers a, b, c

such that $a + b = c$, prove that it is possible for her to eat 100,000 numbers in the first 2011 rows given that she is not restricted to traveling only in the first 2011 rows.

(Here, the $n + 1$ st row of Pascal's triangle consists of entries of the form $\binom{n}{k}$ for integers $0 \leq k \leq n$. Thus, the entry $\binom{n}{k}$ is considered adjacent to the entries $\binom{n-1}{k-1}$, $\binom{n-1}{k}$, $\binom{n}{k-1}$, $\binom{n}{k+1}$, $\binom{n+1}{k}$, $\binom{n+1}{k+1}$.)

Linus Hamilton.

- 4** Consider the infinite grid of lattice points in \mathbb{Z}^3 . Little D and Big Z play a game, where Little D first loses a shoe on an unmunched point in the grid. Then, Big Z munches a shoe-free plane perpendicular to one of the coordinate axes. They continue to alternate turns in this fashion, with Little D's goal to lose a shoe on each of n consecutive lattice points on a line parallel to one of the coordinate axes. Determine all n for which Little D can accomplish his goal.

David Yang.

- 5** Prove there exists a constant c (independent of n) such that for any graph G with $n > 2$ vertices, we can split G into a forest and at most $cf(n)$ disjoint cycles, where

a) $f(n) = n \ln n$;

b) $f(n) = n$.

David Yang.

- 6** Do there exist positive integers k and n such that for any finite graph G with diameter $k + 1$ there exists a set S of at most n vertices such that for any $v \in V(G) \setminus S$, there exists a vertex $u \in S$ of distance at most k from v ?

David Yang.

- 7** Let T be a tree. Prove that there is a constant $c > 0$ (independent of n) such that every graph with n vertices that does not contain a subgraph isomorphic to T has at most cn edges.

David Yang.

– Geometry

- 1** Let $ABCD$ be a convex quadrilateral. Let E, F, G, H be points on segments AB, BC, CD, DA , respectively, and let P be the intersection of EG and FH . Given that quadrilaterals $HAEP$, $EBFP$, $FCGP$, $GDHP$ all have inscribed circles, prove that $ABCD$ also has an inscribed circle.

Evan O'Dorney.

- 2 Let $\omega, \omega_1, \omega_2$ be three mutually tangent circles such that ω_1, ω_2 are externally tangent at P , ω_1, ω are internally tangent at A , and ω, ω_2 are internally tangent at B . Let O, O_1, O_2 be the centers of $\omega, \omega_1, \omega_2$, respectively. Given that X is the foot of the perpendicular from P to AB , prove that $\angle O_1XP = \angle O_2XP$.

David Yang.

- 3 Let ABC be a triangle. Draw circles ω_A, ω_B , and ω_C such that ω_A is tangent to AB and AC , and ω_B and ω_C are defined similarly. Let P_A be the insimilicenter of ω_B and ω_C . Define P_B and P_C similarly. Prove that AP_A, BP_B , and CP_C are concurrent.

Tom Lu.

- 4 Prove that for any convex pentagon $A_1A_2A_3A_4A_5$, there exists a unique pair of points $\{P, Q\}$ (possibly with $P = Q$) such that $\angle PA_iA_{i-1} = \angle A_{i+1}A_iQ$ for $1 \leq i \leq 5$, where indices are taken $(\text{mod } 5)$ and angles are directed $(\text{mod } \pi)$.

Calvin Deng.

– Number Theory

- 1 Prove that $n^3 - n - 3$ is not a perfect square for any integer n .

Calvin Deng.

- 2 Let $p \geq 5$ be a prime. Show that

$$\sum_{k=0}^{(p-1)/2} \binom{p}{k} 3^k \equiv 2^p - 1 \pmod{p^2}.$$

Victor Wang.

- 3 Let $n > 1$ be a fixed positive integer, and call an n -tuple (a_1, a_2, \dots, a_n) of integers greater than 1 *good* if and only if $a_i \mid \left(\frac{a_1 a_2 \cdots a_n}{a_i} - 1 \right)$ for $i = 1, 2, \dots, n$. Prove that there are finitely many good n -tuples.

Mitchell Lee.

- 4 Let $p > 13$ be a prime of the form $2q + 1$, where q is prime. Find the number of ordered pairs of integers (m, n) such that $0 \leq m < n < p - 1$ and

$$3^m + (-12)^m \equiv 3^n + (-12)^n \pmod{p}.$$

Alex Zhu.

The original version asked for the number of solutions to $2^m + 3^n \equiv 2^n + 3^m \pmod{p}$ (still $0 \leq m < n < p - 1$), where p is a Fermat prime.
