

ELMO Shortlist 2012

www.artofproblemsolving.com/community/c4350

by math154

– Algebra

- 1** Let $x_1, x_2, x_3, y_1, y_2, y_3$ be nonzero real numbers satisfying $x_1 + x_2 + x_3 = 0, y_1 + y_2 + y_3 = 0$. Prove that

$$\frac{x_1x_2 + y_1y_2}{\sqrt{(x_1^2 + y_1^2)(x_2^2 + y_2^2)}} + \frac{x_2x_3 + y_2y_3}{\sqrt{(x_2^2 + y_2^2)(x_3^2 + y_3^2)}} + \frac{x_3x_1 + y_3y_1}{\sqrt{(x_3^2 + y_3^2)(x_1^2 + y_1^2)}} \geq -\frac{3}{2}.$$

Ray Li, Max Schindler.

- 2** Let a, b, c be three positive real numbers such that $a \leq b \leq c$ and $a + b + c = 1$. Prove that

$$\frac{a+c}{\sqrt{a^2+c^2}} + \frac{b+c}{\sqrt{b^2+c^2}} + \frac{a+b}{\sqrt{a^2+b^2}} \leq \frac{3\sqrt{6}(b+c)^2}{\sqrt{(a^2+b^2)(b^2+c^2)(c^2+a^2)}}.$$

Owen Goff.

- 3** Prove that any polynomial of the form $1 + a_nx^n + a_{n+1}x^{n+1} + \dots + a_kx^k$ ($k \geq n$) has at least $n - 2$ non-real roots (counting multiplicity), where the a_i ($n \leq i \leq k$) are real and $a_k \neq 0$.

David Yang.

- 4** Let a_0, b_0 be positive integers, and define $a_{i+1} = a_i + \lfloor \sqrt{b_i} \rfloor$ and $b_{i+1} = b_i + \lfloor \sqrt{a_i} \rfloor$ for all $i \geq 0$. Show that there exists a positive integer n such that $a_n = b_n$.

David Yang.

- 5** Prove that if m, n are relatively prime positive integers, $x^m - y^n$ is irreducible in the complex numbers. (A polynomial $P(x, y)$ is irreducible if there do not exist nonconstant polynomials $f(x, y)$ and $g(x, y)$ such that $P(x, y) = f(x, y)g(x, y)$ for all x, y .)

David Yang.

- 6** Let $a, b, c \geq 0$. Show that $(a^2 + 2bc)^{2012} + (b^2 + 2ca)^{2012} + (c^2 + 2ab)^{2012} \leq (a^2 + b^2 + c^2)^{2012} + 2(ab + bc + ca)^{2012}$.

Calvin Deng.

- 7** Let f, g be polynomials with complex coefficients such that $\gcd(\deg f, \deg g) = 1$. Suppose that there exist polynomials $P(x, y)$ and $Q(x, y)$ with complex coefficients such that $f(x) + g(y) = P(x, y)Q(x, y)$. Show that one of P and Q must be constant.

Victor Wang.

- 8** Find all functions $f : \mathbb{Q} \rightarrow \mathbb{R}$ such that $f(x)f(y)f(x+y) = f(xy)(f(x) + f(y))$ for all $x, y \in \mathbb{Q}$.

Sammy Luo and Alex Zhu.

- 9** Let a, b, c be distinct positive real numbers, and let k be a positive integer greater than 3. Show that

$$\left| \frac{a^{k+1}(b-c) + b^{k+1}(c-a) + c^{k+1}(a-b)}{a^k(b-c) + b^k(c-a) + c^k(a-b)} \right| \geq \frac{k+1}{3(k-1)}(a+b+c)$$

and

$$\left| \frac{a^{k+2}(b-c) + b^{k+2}(c-a) + c^{k+2}(a-b)}{a^k(b-c) + b^k(c-a) + c^k(a-b)} \right| \geq \frac{(k+1)(k+2)}{3k(k-1)}(a^2 + b^2 + c^2).$$

Calvin Deng.

- 10** Let $A_1A_2A_3A_4A_5A_6A_7A_8$ be a cyclic octagon. Let B_i be the intersection of A_iA_{i+1} and $A_{i+3}A_{i+4}$. (Take $A_9 = A_1, A_{10} = A_2$, etc.) Prove that B_1, B_2, \dots, B_8 lie on a conic.

David Yang.

– Combinatorics

- 1** Let $n \geq 2$ be a positive integer. Given a sequence (s_i) of n distinct real numbers, define the "class" of the sequence to be the sequence $(a_1, a_2, \dots, a_{n-1})$, where a_i is 1 if $s_{i+1} > s_i$ and -1 otherwise.

Find the smallest integer m such that there exists a sequence (w_i) of length m such that for every possible class of a sequence of length n , there is a subsequence of (w_i) that has that class.

David Yang.

- 2** Determine whether it's possible to cover a K_{2012} with

- a) 1000 K_{1006} 's;
b) 1000 $K_{1006,1006}$'s.

David Yang.

- 3 Find all ordered pairs of positive integers (m, n) for which there exists a set $C = \{c_1, \dots, c_k\}$ ($k \geq 1$) of colors and an assignment of colors to each of the mn unit squares of a $m \times n$ grid such that for every color $c_i \in C$ and unit square S of color c_i , exactly two direct (non-diagonal) neighbors of S have color c_i .

David Yang.

- 4 A tournament on $2k$ vertices contains no 7-cycles. Show that its vertices can be partitioned into two sets, each with size k , such that the edges between vertices of the same set do not determine any 3-cycles.

Calvin Deng.

- 5 Form the infinite graph A by taking the set of primes p congruent to 1 (mod 4), and connecting p and q if they are quadratic residues modulo each other. Do the same for a graph B with the primes 1 (mod 8). Show A and B are isomorphic to each other.

Linus Hamilton.

- 6 Consider a directed graph G with n vertices, where 1-cycles and 2-cycles are permitted. For any set S of vertices, let $N^+(S)$ denote the out-neighborhood of S (i.e. set of successors of S), and define $(N^+)^k(S) = N^+((N^+)^{k-1}(S))$ for $k \geq 2$.

For fixed n , let $f(n)$ denote the maximum possible number of distinct sets of vertices in $\{(N^+)^k(X)\}_{k=1}^\infty$, where X is some subset of $V(G)$. Show that there exists $n > 2012$ such that $f(n) < 1.0001^n$.

Linus Hamilton.

- 7 Consider a graph G with n vertices and at least $n^2/10$ edges. Suppose that each edge is colored in one of c colors such that no two incident edges have the same color. Assume further that no cycles of size 10 have the same set of colors. Prove that there is a constant k such that c is at least $kn^{\frac{8}{5}}$ for any n .

David Yang.

- 8 Consider the equilateral triangular lattice in the complex plane defined by the Eisenstein integers; let the ordered pair (x, y) denote the complex number $x + y\omega$ for $\omega = e^{2\pi i/3}$. We define an ω -chessboard polygon to be a (non self-intersecting) polygon whose sides are situated along lines of the form $x = a$ or $y = b$, where a and b are integers. These lines divide the interior into unit triangles, which are shaded alternately black and white so that adjacent triangles have different colors. To tile an ω -chessboard polygon by lozenges is to exactly cover the polygon by non-overlapping rhombuses consisting of two bordering triangles. Finally, a *tasteful tiling* is one such that for every unit hexagon tiled by three lozenges, each lozenge has a black triangle on its left (defined by clockwise orientation) and a white triangle on its right (so the lozenges are BW, BW, BW in clockwise order).

- a) Prove that if an ω -chessboard polygon can be tiled by lozenges, then it can be done so tastefully.
- b) Prove that such a tasteful tiling is unique.

Victor Wang.

- 9 For a set A of integers, define $f(A) = \{x^2 + xy + y^2 : x, y \in A\}$. Is there a constant c such that for all positive integers n , there exists a set A of size n such that $|f(A)| \leq cn$?

David Yang.

– Geometry

- 1 In acute triangle ABC , let D, E, F denote the feet of the altitudes from A, B, C , respectively, and let ω be the circumcircle of $\triangle AEF$. Let ω_1 and ω_2 be the circles through D tangent to ω at E and F , respectively. Show that ω_1 and ω_2 meet at a point P on BC other than D .

Ray Li.

- 2 In triangle ABC , P is a point on altitude AD . Q, R are the feet of the perpendiculars from P to AB, AC , and QP, RP meet BC at S and T respectively. The circumcircles of BQS and CRT meet QR at X, Y .

a) Prove SX, TY, AD are concurrent at a point Z .

b) Prove Z is on QR iff $Z = H$, where H is the orthocenter of ABC .

Ray Li.

- 3 ABC is a triangle with incenter I . The foot of the perpendicular from I to BC is D , and the foot of the perpendicular from I to AD is P . Prove that $\angle BPD = \angle DPC$.

Alex Zhu.

- 4 Circles Ω and ω are internally tangent at point C . Chord AB of Ω is tangent to ω at E , where E is the midpoint of AB . Another circle, ω_1 is tangent to Ω, ω , and AB at D, Z , and F respectively. Rays CD and AB meet at P . If M is the midpoint of major arc AB , show that $\tan \angle ZEP = \frac{PE}{CM}$.

Ray Li.

- 5 Let ABC be an acute triangle with $AB < AC$, and let D and E be points on side BC such that $BD = CE$ and D lies between B and E . Suppose there exists a point P inside ABC such that $PD \parallel AE$ and $\angle PAB = \angle EAC$. Prove that $\angle PBA = \angle PCA$.

Calvin Deng.

- 6** In $\triangle ABC$, H is the orthocenter, and AD, BE are arbitrary cevians. Let ω_1, ω_2 denote the circles with diameters AD and BE , respectively. HD, HE meet ω_1, ω_2 again at F, G . DE meets ω_1, ω_2 again at P_1, P_2 respectively. FG meets ω_1, ω_2 again at Q_1, Q_2 respectively. P_1H, Q_1H meet ω_1 at R_1, S_1 respectively. P_2H, Q_2H meet ω_2 at R_2, S_2 respectively. Let $P_1Q_1 \cap P_2Q_2 = X$, and $R_1S_1 \cap R_2S_2 = Y$. Prove that X, Y, H are collinear.

Ray Li.

- 7** Let $\triangle ABC$ be an acute triangle with circumcenter O such that $AB < AC$, let Q be the intersection of the external bisector of $\angle A$ with BC , and let P be a point in the interior of $\triangle ABC$ such that $\triangle BPA$ is similar to $\triangle APC$. Show that $\angle QPA + \angle OQB = 90^\circ$.

Alex Zhu.

– Number Theory

- 1** Find all positive integers n such that $4^n + 6^n + 9^n$ is a square.

David Yang, Alex Zhu.

- 2** For positive rational x , if x is written in the form p/q with p, q positive relatively prime integers, define $f(x) = p + q$. For example, $f(1) = 2$.

a) Prove that if $f(x) = f(mx/n)$ for rational x and positive integers m, n , then $f(x)$ divides $|m - n|$.

b) Let n be a positive integer. If all x which satisfy $f(x) = f(2^n x)$ also satisfy $f(x) = 2^n - 1$, find all possible values of n .

Anderson Wang.

- 3** Let $s(k)$ be the number of ways to express k as the sum of distinct 2012^{th} powers, where order does not matter. Show that for every real number c there exists an integer n such that $s(n) > cn$.

Alex Zhu.

- 4** Do there exist positive integers $b, n > 1$ such that when n is expressed in base b , there are more than n distinct permutations of its digits? For example, when $b = 4$ and $n = 18$, $18 = 102_4$, but 102 only has 6 digit arrangements. (Leading zeros are allowed in the permutations.)

Lewis Chen.

- 5** Let $n > 2$ be a positive integer and let p be a prime. Suppose that the nonzero integers are colored in n colors. Let a_1, a_2, \dots, a_n be integers such that for all $1 \leq i \leq n$, $p^i \nmid a_i$ and $p^{i-1} \mid a_i$. In terms of n, p , and $\{a_i\}_{i=1}^n$, determine if there must exist integers x_1, x_2, \dots, x_n of the same color such that $a_1x_1 + a_2x_2 + \dots + a_nx_n = 0$.

Ravi Jagadeesan.

-
- 6 Prove that if a and b are positive integers and $ab > 1$, then

$$\left\lfloor \frac{(a-b)^2 - 1}{ab} \right\rfloor = \left\lfloor \frac{(a-b)^2 - 1}{ab-1} \right\rfloor.$$

Here $\lfloor x \rfloor$ denotes the greatest integer not exceeding x .

Calvin Deng.

- 7 A diabolical combination lock has n dials (each with c possible states), where $n, c > 1$. The dials are initially set to states d_1, d_2, \dots, d_n , where $0 \leq d_i \leq c-1$ for each $1 \leq i \leq n$. Unfortunately, the actual states of the dials (the d_i 's) are concealed, and the initial settings of the dials are also unknown. On a given turn, one may advance each dial by an integer amount c_i ($0 \leq c_i \leq c-1$), so that every dial is now in a state $d'_i \equiv d_i + c_i \pmod{c}$ with $0 \leq d'_i \leq c-1$. After each turn, the lock opens if and only if all of the dials are set to the zero state; otherwise, the lock selects a random integer k and cyclically shifts the d_i 's by k (so that for every i , d_i is replaced by d_{i-k} , where indices are taken modulo n).

Show that the lock can always be opened, regardless of the choices of the initial configuration and the choices of k (which may vary from turn to turn), if and only if n and c are powers of the same prime.

Bobby Shen.

- 8 Fix two positive integers $a, k \geq 2$, and let $f \in \mathbb{Z}[x]$ be a nonconstant polynomial. Suppose that for all sufficiently large positive integers n , there exists a rational number x satisfying $f(x) = f(a^n)^k$. Prove that there exists a polynomial $g \in \mathbb{Q}[x]$ such that $f(g(x)) = f(x)^k$ for all real x .

Victor Wang.

- 9 Are there positive integers m, n such that there exist at least 2012 positive integers x such that both $m - x^2$ and $n - x^2$ are perfect squares?

David Yang.
