

IMC 2012

www.artofproblemsolving.com/community/c4384

by hsiljak, nivotko

Day 1 July 28th

-
- 1** For every positive integer n , let $p(n)$ denote the number of ways to express n as a sum of positive integers. For instance, $p(4) = 5$ because

$$4 = 3 + 1 = 2 + 2 = 2 + 1 + 1 = 1 + 1 + 1.$$

Also define $p(0) = 1$.

Prove that $p(n) - p(n - 1)$ is the number of ways to express n as a sum of integers each of which is strictly greater than 1.

Proposed by Fedor Duzhin, Nanyang Technological University.

-
- 2** Let n be a fixed positive integer. Determine the smallest possible rank of an $n \times n$ matrix that has zeros along the main diagonal and strictly positive real numbers off the main diagonal.

Proposed by Ilya Bogdanov and Grigoriy Chelnokov, MIPT, Moscow.

-
- 3** Given an integer $n > 1$, let S_n be the group of permutations of the numbers $1, 2, 3, \dots, n$. Two players, A and B, play the following game. Taking turns, they select elements (one element at a time) from the group S_n . It is forbidden to select an element that has already been selected. The game ends when the selected elements generate the whole group S_n . The player who made the last move loses the game. The first move is made by A. Which player has a winning strategy?

Proposed by Fedor Petrov, St. Petersburg State University.

-
- 4** Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a continuously differentiable function that satisfies $f'(t) > f(f(t))$ for all $t \in \mathbb{R}$. Prove that $f(f(f(t))) \leq 0$ for all $t \geq 0$.

Proposed by Tom Brta, Charles University, Prague.

-
- 5** Let a be a rational number and let n be a positive integer. Prove that the polynomial $X^{2^n}(X + a)^{2^n} + 1$ is irreducible in the ring $\mathbb{Q}[X]$ of polynomials with rational coefficients.

Proposed by Vincent Jug, cole Polytechnique, Paris.

Day 2 July 29th

1 Consider a polynomial

$$f(x) = x^{2012} + a_{2011}x^{2011} + \cdots + a_1x + a_0.$$

Albert Einstein and Homer Simpson are playing the following game. In turn, they choose one of the coefficients $a_0, a_1, \dots, a_{2011}$ and assign a real value to it. Albert has the first move. Once a value is assigned to a coefficient, it cannot be changed any more. The game ends after all the coefficients have been assigned values.

Homer's goal is to make $f(x)$ divisible by a fixed polynomial $m(x)$ and Albert's goal is to prevent this.

(a) Which of the players has a winning strategy if $m(x) = x - 2012$?

(b) Which of the players has a winning strategy if $m(x) = x^2 + 1$?

Proposed by Fedor Duzhin, Nanyang Technological University.

2 Define the sequence a_0, a_1, \dots inductively by $a_0 = 1$, $a_1 = \frac{1}{2}$, and

$$a_{n+1} = \frac{na_n^2}{1 + (n+1)a_n}, \quad \forall n \geq 1.$$

Show that the series $\sum_{k=0}^{\infty} \frac{a_{k+1}}{a_k}$ converges and determine its value.

Proposed by Christophe Debry, KU Leuven, Belgium.

3 Is the set of positive integers n such that $n! + 1$ divides $(2012n)!$ finite or infinite?

Proposed by Fedor Petrov, St. Petersburg State University.

4 Let $n \geq 2$ be an integer. Find all real numbers a such that there exist real numbers x_1, x_2, \dots, x_n satisfying

$$x_1(1 - x_2) = x_2(1 - x_3) = \cdots = x_n(1 - x_1) = a.$$

Proposed by Walther Janous and Gerhard Kirchner, Innsbruck.

5 Let $c \geq 1$ be a real number. Let G be an Abelian group and let $A \subset G$ be a finite set satisfying $|A + A| \leq c|A|$, where $X + Y := \{x + y | x \in X, y \in Y\}$ and $|Z|$ denotes the cardinality of Z . Prove that

$$|\underbrace{A + A + \cdots + A}_k| \leq c^k |A|$$

for every positive integer k .

Proposed by Przemyslaw Mazur, Jagiellonian University.