

USA Team Selection Test 2013

www.artofproblemsolving.com/community/c4643

by v_Enhance, lyukhson, rrusczyk

– December TST

-
- 1** A social club has $2k + 1$ members, each of whom is fluent in the same k languages. Any pair of members always talk to each other in only one language. Suppose that there were no three members such that they use only one language among them. Let A be the number of three-member subsets such that the three distinct pairs among them use different languages. Find the maximum possible value of A .

-
- 2** Find all triples (x, y, z) of positive integers such that $x \leq y \leq z$ and

$$x^3(y^3 + z^3) = 2012(xyz + 2).$$

-
- 3** Let ABC be a scalene triangle with $\angle BCA = 90^\circ$, and let D be the foot of the altitude from C . Let X be a point in the interior of the segment CD . Let K be the point on the segment AX such that $BK = BC$. Similarly, let L be the point on the segment BX such that $AL = AC$. The circumcircle of triangle DKL intersects segment AB at a second point T (other than D). Prove that $\angle ACT = \angle BCT$.

-
- 4** Let $f : \mathbb{N} \rightarrow \mathbb{N}$ be a function, and let f^m be f applied m times. Suppose that for every $n \in \mathbb{N}$ there exists a $k \in \mathbb{N}$ such that $f^{2k}(n) = n + k$, and let k_n be the smallest such k . Prove that the sequence k_1, k_2, \dots is unbounded.

Proposed by Palmer Mebane, United States

– January TST

-
- 1** Two incongruent triangles ABC and XYZ are called a pair of *pals* if they satisfy the following conditions:
- (a) the two triangles have the same area;
 - (b) let M and W be the respective midpoints of sides BC and YZ . The two sets of lengths $\{AB, AM, AC\}$ and $\{XY, XW, XZ\}$ are identical 3-element sets of pairwise relatively prime integers.

Determine if there are infinitely many pairs of triangles that are pals of each other.

-
- 2** Let ABC be an acute triangle. Circle ω_1 , with diameter AC , intersects side BC at F (other than C). Circle ω_2 , with diameter BC , intersects side AC at E (other than C). Ray AF intersects ω_2

at K and M with $AK < AM$. Ray BE intersects ω_1 at L and N with $BL < BN$. Prove that lines AB, ML, NK are concurrent.

-
- 3** In a table with n rows and $2n$ columns where n is a fixed positive integer, we write either zero or one into each cell so that each row has n zeros and n ones. For $1 \leq k \leq n$ and $1 \leq i \leq n$, we define $a_{k,i}$ so that the i^{th} zero in the k^{th} row is the $a_{k,i}^{\text{th}}$ column. Let \mathcal{F} be the set of such tables with $a_{1,i} \geq a_{2,i} \geq \dots \geq a_{n,i}$ for every i with $1 \leq i \leq n$. We associate another $n \times 2n$ table $f(C)$ from $C \in \mathcal{F}$ as follows: for the k^{th} row of $f(C)$, we write n ones in the columns $a_{n,k} - k + 1, a_{n-1,k} - k + 2, \dots, a_{1,k} - k + n$ (and we write zeros in the other cells in the row).

(a) Show that $f(C) \in \mathcal{F}$.

(b) Show that $f(f(f(f(f(f(C)))))) = C$ for any $C \in \mathcal{F}$.

-
- 4** Determine if there exists a (three-variable) polynomial $P(x, y, z)$ with integer coefficients satisfying the following property: a positive integer n is *not* a perfect square if and only if there is a triple (x, y, z) of positive integers such that $P(x, y, z) = n$.

- https://data.artofproblemsolving.com/images/maa_logo.png These problems are copyright © Mathematical Association of America (<http://maa.org>).
