

AoPS Community

India National Olympiad 1990

www.artofproblemsolving.com/community/c4916 by Leon

1 Given the equation

$$x^4 + px^3 + qx^2 + rx + s = 0$$

has four real, positive roots, prove that (a) $pr - 16s \ge 0$ (b) $q^2 - 36s \ge 0$ with equality in each case holding if and only if the four roots are equal.

2 Determine all non-negative integral pairs (x, y) for which

$$(xy - 7)^2 = x^2 + y^2.$$

3 Let *f* be a function defined on the set of non-negative integers and taking values in the same set. Given that

(a)
$$x - f(x) = 19 \left[\frac{x}{19} \right] - 90 \left[\frac{f(x)}{90} \right]$$
 for all non-negative integers x ;

(b)
$$1900 < f(1990) < 2000$$
,

find the possible values that f(1990) can take. (Notation : here [z] refers to largest integer that is $\leq z$, e.g. [3.1415] = 3).

- **4** Consider the collection of all three-element subsets drawn from the set {1, 2, 3, 4, ..., 299, 300}. Determine the number of those subsets for which the sum of the elements is a multiple of 3.
- **5** Let *a*, *b*, *c* denote the sides of a triangle. Show that the quantity

$$\frac{a}{b+c} + \frac{b}{c+a} + \frac{c}{a+b}$$

must lie between the limits 3/2 and 2. Can equality hold at either limits?

6 Triangle *ABC* is scalene with angle *A* having a measure greater than 90 degrees. Determine the set of points *D* that lie on the extended line *BC*, for which

$$AD| = \sqrt{|BD| \cdot |CD|}$$

where |BD| refers to the (positive) distance between B and D.

AoPS Community

1990 India National Olympiad

7 Let *ABC* be an arbitrary acute angled triangle. For any point *P* lying within the triangle, let *D*, *E*, *F* denote the feet of the perpendiculars from *P* onto the sides *AB*, *BC*, *CA* respectively. Determine the set of all possible positions of the point *P* for which the triangle *DEF* is isosceles.

For which position of P will the triangle DEF become equilateral?

Act of Problem Solving is an ACS WASC Accredited School.

2