

**Baltic Way 2001**

[www.artofproblemsolving.com/community/c5140](http://www.artofproblemsolving.com/community/c5140)

by WakeUp, hasan4444

1 A set of 8 problems was prepared for an examination. Each student was given 3 of them. No two students received more than one common problem. What is the largest possible number of students?

2 Let  $n \geq 2$  be a positive integer. Find whether there exist  $n$  pairwise nonintersecting nonempty subsets of  $\{1, 2, 3, \dots\}$  such that each positive integer can be expressed in a unique way as a sum of at most  $n$  integers, all from different subsets.

3 The numbers  $1, 2, \dots, 49$  are placed in a  $7 \times 7$  array, and the sum of the numbers in each row and in each column is computed. Some of these 14 sums are odd while others are even. Let  $A$  denote the sum of all the odd sums and  $B$  the sum of all even sums. Is it possible that the numbers were placed in the array in such a way that  $A = B$ ?

4 Let  $p$  and  $q$  be two different primes. Prove that

$$\left\lfloor \frac{p}{q} \right\rfloor + \left\lfloor \frac{2p}{q} \right\rfloor + \left\lfloor \frac{3p}{q} \right\rfloor + \dots + \left\lfloor \frac{(q-1)p}{q} \right\rfloor = \frac{1}{2}(p-1)(q-1)$$

5 Let 2001 given points on a circle be coloured either red or green. In one step all points are recoloured simultaneously in the following way: If both direct neighbours of a point  $P$  have the same colour as  $P$ , then the colour of  $P$  remains unchanged, otherwise  $P$  obtains the other colour. Starting with the first colouring  $F_1$ , we obtain the colourings  $F_2, F_3, \dots$  after several recolouring steps. Prove that there is a number  $n_0 \leq 1000$  such that  $F_{n_0} = F_{n_0+2}$ . Is the assertion also true if 1000 is replaced by 999?

6 The points  $A, B, C, D, E$  lie on the circle  $c$  in this order and satisfy  $AB \parallel EC$  and  $AC \parallel ED$ . The line tangent to the circle  $c$  at  $E$  meets the line  $AB$  at  $P$ . The lines  $BD$  and  $EC$  meet at  $Q$ . Prove that  $|AC| = |PQ|$ .

7 Given a parallelogram  $ABCD$ . A circle passing through  $A$  meets the line segments  $AB, AC$  and  $AD$  at inner points  $M, K, N$ , respectively. Prove that

$$|AB| \cdot |AM| + |AD| \cdot |AN| = |AK| \cdot |AC|$$

- 8** Let  $ABCD$  be a convex quadrilateral, and let  $N$  be the midpoint of  $BC$ . Suppose further that  $\angle AND = 135^\circ$ .

Prove that  $|AB| + |CD| + \frac{1}{\sqrt{2}} \cdot |BC| \geq |AD|$ .

- 9** Given a rhombus  $ABCD$ , find the locus of the points  $P$  lying inside the rhombus and satisfying  $\angle APD + \angle BPC = 180^\circ$ .

- 10** In a triangle  $ABC$ , the bisector of  $\angle BAC$  meets the side  $BC$  at the point  $D$ . Knowing that  $|BD| \cdot |CD| = |AD|^2$  and  $\angle ADB = 45^\circ$ , determine the angles of triangle  $ABC$ .

- 11** The real-valued function  $f$  is defined for all positive integers. For any integers  $a > 1, b > 1$  with  $d = \gcd(a, b)$ , we have

$$f(ab) = f(d) \left( f\left(\frac{a}{d}\right) + f\left(\frac{b}{d}\right) \right)$$

Determine all possible values of  $f(2001)$ .

- 12** Let  $a_1, a_2, \dots, a_n$  be positive real numbers such that  $\sum_{i=1}^n a_i^3 = 3$  and  $\sum_{i=1}^n a_i^5 = 5$ . Prove that  $\sum_{i=1}^n a_i > \frac{3}{2}$ .

- 13** Let  $a_0, a_1, a_2, \dots$  be a sequence of real numbers satisfying  $a_0 = 1$  and  $a_n = a_{\lfloor 7n/9 \rfloor} + a_{\lfloor n/9 \rfloor}$  for  $n = 1, 2, \dots$

Prove that there exists a positive integer  $k$  with  $a_k < \frac{k}{2001!}$ .

- 14** There are  $2n$  cards. On each card some real number  $x$ , ( $1 \leq x \leq 2n$ ), is written (there can be different numbers on different cards). Prove that the cards can be divided into two heaps with sums  $s_1$  and  $s_2$  so that  $\frac{n}{n+1} \leq \frac{s_1}{s_2} \leq 1$ .

- 15** Let  $a_0, a_1, a_2, \dots$  be a sequence of positive real numbers satisfying  $i \cdot a_i \geq (i+1) \cdot a_{i+1}$  for  $i = 1, 2, \dots$ . Furthermore, let  $x$  and  $y$  be positive reals, and let  $b_i = xa_i + ya_{i-1}$  for  $i = 1, 2, \dots$ . Prove that the inequality  $i \cdot b_i \geq (i+1) \cdot b_{i+1}$  holds for all integers  $i \geq 2$ .

- 16** Let  $f$  be a real-valued function defined on the positive integers satisfying the following condition: For all  $n > 1$  there exists a prime divisor  $p$  of  $n$  such that  $f(n) = f\left(\frac{n}{p}\right) - f(p)$ . Given that  $f(2001) = 1$ , what is the value of  $f(2002)$ ?

- 17** Let  $n$  be a positive integer. Prove that at least  $2^{n-1} + n$  numbers can be chosen from the set  $\{1, 2, 3, \dots, 2^n\}$  such that for any two different chosen numbers  $x$  and  $y$ ,  $x + y$  is not a divisor of  $x \cdot y$ .

- 18** Let  $a$  be an odd integer. Prove that  $a^{2^m} + 2^{2^m}$  and  $a^{2^n} + 2^{2^n}$  are relatively prime for all positive integers  $n$  and  $m$  with  $n \neq m$ .

---

**19** What is the smallest positive odd integer having the same number of positive divisors as 360?

---

**20** From a sequence of integers  $(a, b, c, d)$  each of the sequences

$$(c, d, a, b), \quad (b, a, d, c), \quad (a + nc, b + nd, c, d), \quad (a + nb, b, c + nd, d)$$

for arbitrary integer  $n$  can be obtained by one step. Is it possible to obtain  $(3, 4, 5, 7)$  from  $(1, 2, 3, 4)$  through a sequence of such steps?

---