

AoPS Community

Baltic Way 2002

www.artofproblemsolving.com/community/c5141 by WakeUp, srinath.r, MJ GEO, stergiu

1 Solve the system of simultaneous equations

$$\begin{cases} a^3 + 3ab^2 + 3ac^2 - 6abc = 1\\ b^3 + 3ba^2 + 3bc^2 - 6abc = 1\\ c^3 + 3ca^2 + 3cb^2 - 6abc = 1 \end{cases}$$

in real numbers.

2 Let a, b, c, d be real numbers such that

$$a + b + c + d = -2$$

$$ab + ac + ad + bc + bd + cd = 0$$

Prove that at least one of the numbers a, b, c, d is not greater than -1.

3 Find all sequences $0 \le a_0 \le a_1 \le a_2 \le \dots$ of real numbers such that

$$a_{m^2+n^2} = a_m^2 + a_n^2$$

for all integers $m, n \geq 0$.

4 Let n be a positive integer. Prove that

$$\sum_{i=1}^{n} x_i (1 - x_i)^2 \le \left(1 - \frac{1}{n}\right)^2$$

for all nonnegative real numbers x_1, x_2, \ldots, x_n such that $x_1 + x_2 + \ldots + x_n = 1$.

5 Find all pairs (a, b) of positive rational numbers such that

$$\sqrt{a} + \sqrt{b} = \sqrt{2 + \sqrt{3}}.$$

The following solitaire game is played on an $m \times n$ rectangular board, $m,n \geq 2$, divided into unit squares. First, a rook is placed on some square. At each move, the rook can be moved an arbitrary number of squares horizontally or vertically, with the extra condition that each move has to be made in the 90° clockwise direction compared to the previous one (e.g. after a move

to the left, the next one has to be done upwards, the next one to the right etc). For which values of m and n is it possible that the rook visits every square of the board exactly once and returns to the first square? (The rook is considered to visit only those squares it stops on, and not the ones it steps over.)

- 7 We draw n convex quadrilaterals in the plane. They divide the plane into regions (one of the regions is infinite). Determine the maximal possible number of these regions.
- Let P be a set of $n \ge 3$ points in the plane, no three of which are on a line. How many possibilities are there to choose a set T of $\binom{n-1}{2}$ triangles, whose vertices are all in P, such that each triangle in T has a side that is not a side of any other triangle in T?
- Two magicians show the following trick. The first magician goes out of the room. The second magician takes a deck of 100 cards labelled by numbers 1, 2, ..., 100 and asks three spectators to choose in turn one card each. The second magician sees what card each spectator has taken. Then he adds one more card from the rest of the deck. Spectators shuffle these 4 cards, call the first magician and give him these 4 cards. The first magician looks at the 4 cards and guesses what card was chosen by the first spectator, what card by the second and what card by the third. Prove that the magicians can perform this trick.
- Let N be a positive integer. Two persons play the following game. The first player writes a list of positive integers not greater than 25, not necessarily different, such that their sum is at least 200. The second player wins if he can select some of these numbers so that their sum S satisfies the condition $200 N \le S \le 200 + N$. What is the smallest value of N for which the second player has a winning strategy?
- Let n be a positive integer. Consider n points in the plane such that no three of them are collinear and no two of the distances between them are equal. One by one, we connect each point to the two points nearest to it by line segments (if there are already other line segments drawn to this point, we do not erase these). Prove that there is no point from which line segments will be drawn to more than 11 points.
- A set S of four distinct points is given in the plane. It is known that for any point $X \in S$ the remaining points can be denoted by Y, Z and W so that |XY| = |XZ| + |XW| Prove that all four points lie on a line.
- Let ABC be an acute triangle with $\angle BAC > \angle BCA$, and let D be a point on side AC such that |AB| = |BD|. Furthermore, let F be a point on the circumcircle of triangle ABC such that line FD is perpendicular to side BC and points F, B lie on different sides of line AC. Prove that line FB is perpendicular to side AC.
- Let L, M and N be points on sides AC, AB and BC of triangle ABC, respectively, such that BL is the bisector of angle ABC and segments AN, BL and CM have a common point. Prove

that if $\angle ALB = \angle MNB$ then $\angle LNM = 90^{\circ}$.

A spider and a fly are sitting on a cube. The fly wants to maximize the shortest path to the 15 spider along the surface of the cube. Is it necessarily best for the fly to be at the point opposite to the spider?

(Opposite means symmetric with respect to the centre of the cube.)

16 Find all nonnegative integers m such that

$$a_m = (2^{2m+1})^2 + 1$$

is divisible by at most two different primes.

17 Show that the sequence

$$\binom{2002}{2002}$$
, $\binom{2003}{2002}$, $\binom{2004}{2002}$, ...

considred modulo 2002, is periodic.

- Find all integers n > 1 such that any prime divisor of $n^6 1$ is a divisor of $(n^3 1)(n^2 1)$. 18
- 19 Let n be a positive integer. Prove that the equation

$$x + y + \frac{1}{x} + \frac{1}{y} = 3n$$

does not have solutions in positive rational numbers.

Does there exist an infinite non-constant arithmetic progression, each term of which is of the 20 form a^b , where a and b are positive integers with b > 2?