

**Iran Team Selection Test 2007**

[www.artofproblemsolving.com/community/c5383](http://www.artofproblemsolving.com/community/c5383)

by Omid Hatami, pohoatza

**Day 1**

- 
- 1 In triangle  $ABC$ ,  $M$  is midpoint of  $AC$ , and  $D$  is a point on  $BC$  such that  $DB = DM$ . We know that  $2BC^2 - AC^2 = AB \cdot AC$ . Prove that

$$BD \cdot DC = \frac{AC^2 \cdot AB}{2(AB + AC)}$$

- 
- 2 Let  $A$  be the largest subset of  $\{1, \dots, n\}$  such that for each  $x \in A$ ,  $x$  divides at most one other element in  $A$ . Prove that

$$\frac{2n}{3} \leq |A| \leq \left\lceil \frac{3n}{4} \right\rceil.$$

- 
- 3 Find all solutions of the following functional equation:

$$f(x^2 + y + f(y)) = 2y + f(x)^2.$$

---

**Day 2**

- 
- 1 In an isosceles right-angled triangle shaped billiards table, a ball starts moving from one of the vertices adjacent to hypotenuse. When it reaches to one side then it will reflect its path. Prove that if we reach to a vertex then it is not the vertex at initial position  
*By Sam Nariman*

- 
- 2 Find all monic polynomials  $f(x)$  in  $\mathbb{Z}[x]$  such that  $f(\mathbb{Z})$  is closed under multiplication.  
*By Mohsen Jamali*

- 
- 3 Let  $\omega$  be incircle of  $ABC$ .  $P$  and  $Q$  are on  $AB$  and  $AC$ , such that  $PQ$  is parallel to  $BC$  and is tangent to  $\omega$ .  $AB, AC$  touch  $\omega$  at  $F, E$ . Prove that if  $M$  is midpoint of  $PQ$ , and  $T$  is intersection point of  $EF$  and  $BC$ , then  $TM$  is tangent to  $\omega$ .  
*By Ali Khezeli*

---

**Day 3**

---

- 1 Does there exist a sequence  $a_0, a_1, a_2, \dots$  in  $\mathbb{N}$ , such that for each  $i \neq j$ ,  $(a_i, a_j) = 1$ , and for each  $n$ , the polynomial  $\sum_{i=0}^n a_i x^i$  is irreducible in  $\mathbb{Z}[x]$ ?

By Omid Hatami

- 2 Suppose  $n$  lines in plane are such that no two are parallel and no three are concurrent. For each two lines their angle is a real number in  $[0, \frac{\pi}{2}]$ . Find the largest value of the sum of the  $\binom{n}{2}$  angles between line.

By Aliakbar Daemi

- 3  $O$  is a point inside triangle  $ABC$  such that  $OA = OB + OC$ . Suppose  $B', C'$  be midpoints of arcs  $AOC$  and  $AOB$ . Prove that circumcircles  $COC'$  and  $BOB'$  are tangent to each other.

#### Day 4

- 1 Find all polynomials of degree 3, such that for each  $x, y \geq 0$ :

$$p(x + y) \geq p(x) + p(y)$$

- 2 Triangle  $ABC$  is isosceles ( $AB = AC$ ). From  $A$ , we draw a line  $\ell$  parallel to  $BC$ .  $P, Q$  are on perpendicular bisectors of  $AB, AC$  such that  $PQ \perp BC$ .  $M, N$  are points on  $\ell$  such that angles  $\angle APM$  and  $\angle AQN$  are  $\frac{\pi}{2}$ . Prove that

$$\frac{1}{AM} + \frac{1}{AN} \leq \frac{2}{AB}$$

- 3 Let  $P$  be a point in a square whose side are mirror. A ray of light comes from  $P$  and with slope  $\alpha$ . We know that this ray of light never arrives to a vertex. We make an infinite sequence of 0, 1. After each contact of light ray with a horizontal side, we put 0, and after each contact with a vertical side, we put 1. For each  $n \geq 1$ , let  $B_n$  be set of all blocks of length  $n$ , in this sequence.
- a) Prove that  $B_n$  does not depend on location of  $P$ .
- b) Prove that if  $\frac{\alpha}{\pi}$  is irrational, then  $|B_n| = n + 1$ .