

USA TSTST 2019

www.artofproblemsolving.com/community/c898138

by MarkBcc168, pinetree1, tastymath75025

– Day 1

-
- 1** Find all binary operations $\diamond : \mathbb{R}_{>0} \times \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$ (meaning \diamond takes pairs of positive real numbers to positive real numbers) such that for any real numbers $a, b, c > 0$,
- the equation $a \diamond (b \diamond c) = (a \diamond b) \cdot c$ holds; and
 - if $a \geq 1$ then $a \diamond a \geq 1$.

Evan Chen

-
- 2** Let ABC be an acute triangle with circumcircle Ω and orthocenter H . Points D and E lie on segments AB and AC respectively, such that $AD = AE$. The lines through B and C parallel to \overline{DE} intersect Ω again at P and Q , respectively. Denote by ω the circumcircle of $\triangle ADE$.

- Show that lines PE and QD meet on ω .
- Prove that if ω passes through H , then lines PD and QE meet on ω as well.

Merlijn Staps

-
- 3** On an infinite square grid we place finitely many *cars*, which each occupy a single cell and face in one of the four cardinal directions. Cars may never occupy the same cell. It is given that the cell immediately in front of each car is empty, and moreover no two cars face towards each other (no right-facing car is to the left of a left-facing car within a row, etc.). In a *move*, one chooses a car and shifts it one cell forward to a vacant cell. Prove that there exists an infinite sequence of valid moves using each car infinitely many times.

Nikolai Beluhov

– Day 2

-
- 4** Consider coins with positive real denominations not exceeding 1. Find the smallest $C > 0$ such that the following holds: if we have any 100 such coins with total value 50, then we can always split them into two stacks of 50 coins each such that the absolute difference between the total values of the two stacks is at most C .

Merlijn Staps

-
- 5** Let ABC be an acute triangle with orthocenter H and circumcircle Γ . A line through H intersects segments AB and AC at E and F , respectively. Let K be the circumcenter of $\triangle AEF$,

and suppose line AK intersects Γ again at a point D . Prove that line HK and the line through D perpendicular to \overline{BC} meet on Γ .

Gunmay Handa

-
- 6** Suppose P is a polynomial with integer coefficients such that for every positive integer n , the sum of the decimal digits of $|P(n)|$ is not a Fibonacci number. Must P be constant? (A *Fibonacci number* is an element of the sequence F_0, F_1, \dots defined recursively by $F_0 = 0, F_1 = 1$, and $F_{k+2} = F_{k+1} + F_k$ for $k \geq 0$.)

Nikolai Beluhov

– Day 3

-
- 7** Let $f : \mathbb{Z} \rightarrow \{1, 2, \dots, 10^{100}\}$ be a function satisfying

$$\gcd(f(x), f(y)) = \gcd(f(x), x - y)$$

for all integers x and y . Show that there exist positive integers m and n such that $f(x) = \gcd(m + x, n)$ for all integers x .

Ankan Bhattacharya

-
- 8** Let S be a set of 16 points in the plane, no three collinear. Let $\chi(S)$ denote the number of ways to draw 8 lines with endpoints in S , such that no two drawn segments intersect, even at endpoints. Find the smallest possible value of $\chi(S)$ across all such S .

Ankan Bhattacharya

-
- 9** Let ABC be a triangle with incenter I . Points K and L are chosen on segment BC such that the incircles of $\triangle ABK$ and $\triangle ABL$ are tangent at P , and the incircles of $\triangle ACK$ and $\triangle ACL$ are tangent at Q . Prove that $IP = IQ$.

Ankan Bhattacharya
