

AoPS Community

Math Prize for Girls Olympiad 2015

www.artofproblemsolving.com/community/c952042 by Ravi B

1 Prove that every positive integer has a unique representation in the form

$$\sum_{i=0}^k d_i 2^i \,,$$

where k is a nonnegative integer and each d_i is either 1 or 2. (This representation is similar to usual binary notation except that the digits are 1 and 2, not 0 and 1.)

- **2** A tetrahedron *T* is inside a cube *C*. Prove that the volume of *T* is at most one-third the volume of *C*.
- **3** Let *f* be the cubic polynomial

$$f(x) = x^3 + bx^2 + cx + d,$$

where *b*, *c*, and *d* are real numbers. Let $x_1, x_2, ..., x_n$ be nonnegative numbers, and let *m* be their average. Suppose that $m \ge -\frac{b}{2}$. Prove that

$$\sum_{i=1}^{n} f(x_i) \ge n f(m).$$

An 8-by-8 square is divided into 64 unit squares in the usual way. Each unit square is colored black or white. The number of black unit squares is even. We can take two adjacent unit squares (forming a 1-by-2 or 2-by-1 rectangle), and flip their colors: black becomes white and white becomes black. We call this operation a *step*. If *C* is the original coloring, let S(C) be the least number of steps required to make all the unit squares black. Find with proof the greatest possible value of S(C).

AoPS Online AoPS Academy AoPS Catery

© 2019 AoPS Incorporated 1

Art of Problem Solving is an ACS WASC Accredited School.